

A NEW QUASI-LOCAL MASS AND POSITIVITY

XIAO ZHANG

ABSTRACT. A new definition of quasi-local mass is proposed and its positivity is proved under certain conditions.

1. INTRODUCTION

Recently, Kijowski [12], Liu and Yau [13] propose a definition of quasi-local mass for any spacelike, topological 2-sphere with positive intrinsic (or Gauss) curvature. In particular, Liu and Yau [13, 14] are able to prove its positivity. Liu and Yau's definition arises naturally from calculations in the recent work of Yau [22] on black holes as a candidate for positivity. They use the original idea of Schoen and Yau [18, 19] for the proof of the positive mass conjecture to solve Jang's equation and reduce to the Riemannian version of the positivity proved earlier by Shi and Tam [17].

As it is well-known, there have been many attempts to define a quasi-local mass or energy associated to a spacelike 2-surface. Certain criterions which a definition should have were posed by Christodoulou and Yau [6], and Liu and Yau [14]. However, no known definition can satisfy all these criterions.

The definition of Kijowski, Liu and Yau belongs to the class of definitions initiated by Brown and York [2, 3] for a spacelike 2-surface which bounds a compact spacelike hypersurface in a time orientable space-time. Similar derivation was also obtained by Hawking and Horowitz [9]. We refer to [15] for a short exploiting of the idea of Brown and York. When the spacelike 2-sphere bounds a totally geodesic spacelike hypersurface, the definition of Kijowski, Liu and Yau reduces to the quasi-local mass of Brown and York whose positivity was proved in [17] for topological 2-sphere with positive Gauss curvature and positive mean curvature.

The quasi-local energy-momentum of Brown and York has some desirable properties. It approaches to Arnowitt-Deser-Misner (ADM)

2000 *Mathematics Subject Classification.* 53C27, 53C50, 83C60.

Key words and phrases. General relativity, quasi-local mass, positivity.

Partially supported by NSF of China(10421001), NKBRPC(2006CB805905) and the Innovation Project of Chinese Academy of Sciences.

energy-momentum in a suitable limit [2, 4]. However, the decreasing monotonicity property indicates that the larger region contains less mass (see Lemma 4.2 [17], also [14] for the computation of the round sphere in time slices in the Schwarzschild spacetime). And this breaks certain physical intuition.

To obtain an increasing monotonicity quasi-local mass of Brown and York's type, we propose a new definition by choosing certain spinor norm as lapse function. The idea of using spinor norm coupled with the mean curvature to define the quasi local mass and momentum was told by Yau in summer 2005, which was used in [20]. If a spacelike topological 2-sphere bounds a smooth spacelike hypersurface (initial data set) in a spacetime whose mean curvature does not change sign, the mean curvature of this topological sphere satisfies apparent horizon conditions and the topological sphere can be smoothly isometrically embedded into Euclidean 3-space whose image has nonnegative mean curvature, then the positivity holds. Moreover, if the initial data set has the zero second fundamental form, the vanishing quasi-local mass implies the spacelike topological 2-sphere bounds a domain in Euclidean 3-space. Here, we don't assume the mean curvature of spacelike 2-sphere is strictly positive as [17, 13, 14]. Thus our theorem of positivity can apply to apparent horizons, in particular, to the round sphere in time slices in the Schwarzschild spacetime. Indeed, we compute our quasi-local mass in this case and find that it is an increasing function of the radius of the round sphere.

2. PRELIMINARY

Let $(N, \tilde{g})$ be a 4-dimensional spacetime which satisfies the Einstein field equations

$$\widetilde{Ric} - \frac{\tilde{R}}{2} \tilde{g} = T,$$

where $\widetilde{Ric}$, $\tilde{R}$ are Ricci curvature tensor, scalar curvature of $\tilde{g}$ respectively, and T is the energy-momentum tensor. Let (M, g, p) be a smooth *initial data set*, where M is a spacelike (orientable) hypersurface of N with the smooth induced Riemannian metric g and second fundamental form p . We will introduce briefly the Dirac-Witten operator along M . We refer to [21, 16, 23] for the basic materials.

Fix a point $p \in M$ and an orthonormal basis $\{e_\alpha\}$ of $T_p N$ with e_0 normal and e_i tangent to M ($1 \leq i \leq 3$). Let ω^β_α be the connection 1-form

$$\tilde{\nabla} e_\alpha = e_\beta \otimes \omega^\beta_\alpha. \quad (2.1)$$

Denote

$$\omega_{\alpha\beta} = g_{\alpha\gamma}\omega^\gamma_\beta.$$

Since

$$\begin{aligned} 0 &= d(e_\alpha, e_\beta) \\ &= (\tilde{\nabla} e_\alpha, e_\beta) + (e_\alpha, \tilde{\nabla} e_\beta) \\ &= (e_\gamma \otimes \omega^\gamma_\alpha, e_\beta) + (e_\alpha, e_\delta \otimes \omega^\delta_\beta), \end{aligned}$$

we have

$$\omega_{\alpha\beta} = -\omega_{\beta\alpha}.$$

Thus

$$\omega^0_i = -\omega_{0i} = \omega_{i0} = \omega^i_0.$$

Note that for any vector X

$$(e_\alpha \wedge e_\beta)(X) = \tilde{g}(e_\alpha, X)e_\beta - \tilde{g}(e_\beta, X)e_\alpha.$$

This implies

$$\begin{aligned} (e_0 \wedge e_j)(e_0, e_1, e_2, e_3) &= (e_0, e_1, e_2, e_3)E_{0j}, \\ (e_i \wedge e_j)(e_0, e_1, e_2, e_3) &= (e_0, e_1, e_2, e_3)E_{ij} \end{aligned}$$

where $E_{0j} = (a^0_{\alpha\beta})$ with $a^0_{0j} = a^0_{j0} = -1$ for $j \neq 0$, *others* = 0; $E_{ij} = (a_{\alpha\beta})$ with $a_{ij} = -a_{ji} = -1$ for $i < j \neq 0$, *others* = 0.

Denote the 3-vector Ω_0 and 3×3 anti-symmetric matrix Ω by

$$\begin{aligned} \Omega_0 &= (\omega_{01}, \omega_{02}, \omega_{03}), \\ \Omega &= (\omega_{ij}). \end{aligned}$$

Now (2.1) gives that

$$\begin{aligned} \tilde{\nabla} e_0 &= -e_i \otimes \omega_{0i}, \\ \tilde{\nabla} e_i &= -e_j \otimes \omega_{ij} + e_0 \otimes \omega_{i0}, \end{aligned}$$

i.e.,

$$\tilde{\nabla}(e_0, e_1, e_2, e_3) = (e_0, e_1, e_2, e_3) \otimes \begin{pmatrix} 0 & -\Omega_0 \\ -\Omega_0^t & \Omega \end{pmatrix}.$$

This induces a connection on the principal $SO(3, 1)$ bundle with the connection 1-form

$$\omega = -\frac{1}{2}\omega_{ij}e_i \wedge e_j + \omega_{0i}e_0 \wedge e_i.$$

Denote by $\mathbb{S}$ the (local) spinor bundle of N . Since M is spin, $\mathbb{S}$ exists globally over M . This spinor bundle $\mathbb{S}$ is the hypersurface spinor bundle of M . Let $\tilde{\nabla}$ and $\bar{\nabla}$ be the Levi-Civita connections of $\tilde{g}$ and g respectively, the same symbols are used to denote their lifts to the

hypersurface spinor bundle. There exists a Hermitian inner product $(\cdot, \cdot)$ on $\mathbb{S}$ along M which is compatible with the spin connection $\tilde{\nabla}$. The Clifford multiplication of any vector $\tilde{X}$ of N is symmetric with respect to this inner product. However, this inner product is not positive definite and there exists a positive definite Hermitian inner product defined by $\langle \cdot, \cdot \rangle = (e_0 \cdot, \cdot)$ on $\mathbb{S}$ along M .

Denote $\{\sigma_a\}$ the orthonormal basis on the hypersurface spinor bundle $\mathbb{S}$. Then the (local) spin connection of N is lifted from the principal $SO(3, 1)$ connection as

$$\tilde{\nabla}\sigma_a = -\frac{1}{4}\omega_{ij} \otimes e_i \cdot e_j \cdot \sigma_a + \frac{1}{2}\omega_{0i} \otimes e_0 \cdot e_i \cdot \sigma_a.$$

For spinor $\phi = u^a \sigma_a$, vector $\tilde{X} \in \Gamma(TN)$, we have

$$\begin{aligned} \tilde{\nabla}_{\tilde{X}}\phi &= du^a(\tilde{X})\sigma_a - \frac{u^a}{4}\omega_{ij}(\tilde{X})e_i \cdot e_j \cdot \sigma_a \\ &\quad + \frac{u^a}{2}\omega_{i0}(\tilde{X})e_i \cdot e_0 \cdot \sigma_a. \end{aligned}$$

Therefore, for $X \in \Gamma(TM)$,

$$\tilde{\nabla}_X\phi = \overline{\nabla}_X\phi + \frac{1}{2}\omega_{0i}(X)e_0 \cdot e_i \cdot \phi. \quad (2.2)$$

Define the second fundamental form of the initial data set

$$p_{ij} = \tilde{g}(\tilde{\nabla}_i e_0, e_j) = \omega_{j0}(e_i).$$

Then

$$\tilde{\nabla}_i = \overline{\nabla}_i - \frac{1}{2}p_{ij}e_0 \cdot e_j \cdot. \quad (2.3)$$

This implies that the spinor connection $\overline{\nabla}$ is compatible with the positive definite inner product $\langle \cdot, \cdot \rangle$.

The Dirac-Witten operator along M is defined by

$$\tilde{D} = e_i \cdot \tilde{\nabla}_i.$$

The Dirac operator of M but acting on $\mathbb{S}$ is defined by

$$\overline{D} = e_i \cdot \overline{\nabla}_i.$$

By (2.3), we obtain

$$\tilde{D} = \overline{D} - \frac{tr_g(p)}{2}e_0 \cdot. \quad (2.4)$$

The operator $\tilde{D}$ is formally self-adjoint with respect to $\langle \cdot, \cdot \rangle$ and its square is given by the following Weitzenböck type formula

$$\tilde{D}^2 = \tilde{\nabla}^* \tilde{\nabla} + \mathcal{T} \quad (2.5)$$

where

$$\mathcal{T} = \frac{1}{2}(T_{00} + T_{0i}e_0 \cdot e_i),$$

and $\tilde{\nabla}^*$ is the formal adjoint of $\tilde{\nabla}$ with respect to $\langle \cdot, \cdot \rangle$. If the space-time satisfies the *dominant energy condition*, then $\mathcal{T}$ is a nonnegative operator.

Suppose that M has boundary Σ which has finitely many connected components $\Sigma_1, \dots, \Sigma_l$, each of which is a topological 2-sphere, endowed with its induced Riemannian and spin structures. Fix a point $p \in \Sigma$ and an orthonormal basis $\{e_i\}$ of $T_p M$ with $e_r = e_1$ the outward normal to Σ and e_a tangent to Σ for $2 \leq a \leq 3$. Let

$$h_{ab} = \langle \bar{\nabla}_a e_r, e_b \rangle$$

be the component of the second fundamental form of Σ . Let $H = \text{tr}(h)$ be its mean curvature. Σ is a *future/past apparent horizon* if

$$H \mp \text{tr}(p|_\Sigma) \geq 0 \quad (2.6)$$

holds on Σ . When Σ has multi-components, we require that (2.6) holds (with the same sign) on each Σ_i .

Denote by ∇ the lift of the Levi-Civita connection of Σ to the spinor bundle $\mathbb{S}|_\Sigma$. Then

$$\bar{\nabla}_a = \nabla_a + \frac{1}{2}h_{ab}e_r \cdot e_b \cdot \cdot \quad (2.7)$$

The connection ∇ is also compatible with the metric $\langle \cdot, \cdot \rangle$ on the boundary Σ . Let

$$D = e_a \cdot \nabla_a$$

be the Dirac operator of Σ but acting on $\mathbb{S}|_\Sigma$. From (2.3), (2.7), we obtain

$$\tilde{\nabla}_a = \nabla_a + \frac{1}{2}h_{ab}e_r \cdot e_b \cdot - \frac{1}{2}p_{aj}e_0 \cdot e_j \cdot \cdot \quad (2.8)$$

Therefore (2.5) gives rise to

$$\begin{aligned} \int_M |\tilde{\nabla}\phi|^2 + \langle \phi, \mathcal{T}\phi \rangle - |\tilde{D}\phi|^2 &= \int_\Sigma \langle \phi, (e_r \cdot D - \frac{H}{2})\phi \rangle \\ &\quad + \frac{1}{2} \int_\Sigma \langle \phi, \text{tr}(p|_\Sigma)e_0 \cdot e_r \cdot \phi \rangle \\ &\quad - \frac{1}{2} \int_\Sigma \langle \phi, p_{ar}e_0 \cdot e_a \cdot \phi \rangle. \end{aligned} \quad (2.9)$$

We will prove the following existences. Let

$$P_{\pm} = \frac{1}{2}(Id \pm e_0 \cdot e_r \cdot)$$

be the projective operators on $\mathbb{S}|_{\Sigma}$. The local boundary conditions

$$P_{\pm}\phi|_{\Sigma} = 0$$

ensure that $\tilde{D}$ is a formally self-adjoint elliptic operator [8].

Proposition 1. *Let $(N, \tilde{g})$ be a spacetime which satisfies the dominant energy condition. Let (M, g, p) be a smooth initial data with the boundary Σ which has finitely many multi-components Σ_i , each of which is topological 2-sphere.*

(i) *If $tr_g(p) \geq 0$ and Σ is a past apparent horizon, then the following Dirac-Witten equation has a unique smooth solution $\phi \in \Gamma(\mathbb{S})$*

$$\begin{cases} \tilde{D}\phi &= 0 & \text{in} & M \\ P_+\phi &= P_+\phi_0 & \text{on} & \Sigma_{i_0} \\ P_+\phi &= 0 & \text{on} & \Sigma_i \ (i \neq i_0) \end{cases} \quad (2.10)$$

for any given $\phi_0 \in \Gamma(\mathbb{S}|_{\Sigma})$ and for fixed i_0 ;

(ii) *If $tr_g(p) \leq 0$ and Σ is a future apparent horizon, then the following Dirac-Witten equation has a unique smooth solution $\phi \in \Gamma(\mathbb{S})$*

$$\begin{cases} \tilde{D}\phi &= 0 & \text{in} & M \\ P_-\phi &= P_-\phi_0 & \text{on} & \Sigma_{i_0} \\ P_-\phi &= 0 & \text{on} & \Sigma_i \ (i \neq i_0) \end{cases} \quad (2.11)$$

for any given $\phi_0 \in \Gamma(\mathbb{S}|_{\Sigma})$ and for fixed i_0 .

Proof : To establish the existences of (2.10) and (2.11), we only need to show that any solution $\tilde{D}\phi = 0$ is trivial when $\phi_0 = 0$. By (2.4), we obtain

$$\tilde{D}(e_0 \cdot \phi) = -tr_g(p)\phi.$$

Therefore,

$$\begin{aligned} \int_{\Sigma} \langle e_r \cdot \phi, e_0 \cdot \phi \rangle &= \int_M \langle \tilde{D}\phi, e_0 \cdot \phi \rangle - \langle \phi, \tilde{D}(e_0 \cdot \phi) \rangle \\ &= \int_M tr_g(p)|\phi|^2. \end{aligned}$$

If Σ is a past apparent horizon and $tr_g(p) \geq 0$, or Σ is a future apparent horizon and $tr_g(p) \leq 0$, then $\phi|_{\Sigma} = 0$. Note that, under the local

boundary conditions, (2.9) becomes

$$\int_M |\tilde{\nabla}\phi|^2 + \langle \phi, \mathcal{T} \cdot \phi \rangle - |\tilde{D}\phi|^2 = -\frac{1}{2} \int_\Sigma [H \pm \text{tr}(p|_\Sigma)] |\phi|^2. \quad (2.12)$$

This implies $\tilde{\nabla}_i \phi = 0$ on M . Thus

$$|d|\phi|^2| \leq 2|\tilde{\nabla}\phi||\phi| \leq |p||\phi|^2 \leq C_p |\phi|^2,$$

where $C_p = \max|p|$. If there exists a point $x_0 \in M$ such that $\phi(x_0) \neq 0$, then

$$|\phi|(x) \geq |\phi|(x_0)e^{-C_p \rho_{x_0}(x)}.$$

Taking x to the boundary will give rise to a contradiction. Therefore $\phi = 0$ on M and the existences of (2.10) and (2.11) follow. Q.E.D.

3. DEFINITION AND POSITIVITY

Let $(N, \tilde{g})$ be a spacetime which satisfies the dominant energy condition. Let (M, g, p) be a smooth initial data set with the boundary Σ which has finitely many multi-components Σ_i , each of which is topological 2-sphere. Let e_r be the unit vector outward normal to Σ , h_{ij} and H be the second fundamental form and the mean curvature of Σ respectively.

Suppose that some Σ_{i_0} can be smoothly isometrically embedded into $\mathbb{R}^3$ in the Minkowski spacetime $\mathbb{R}^{3,1}$ and denote by $\aleph$ the isometric embedding. (It exists if Σ_{i_0} has positive Gauss curvature.) Let $\check{\Sigma}$ be the image of Σ under the map $\aleph$. Let $\check{e}_r$ the unit vector outward normal to $\check{\Sigma}$ and $\check{h}_{ij}$, $\check{H}$ are the second fundamental forms, the mean curvature of $\check{\Sigma}$ respectively. Denote by

$$H_0 = \check{H} \circ \aleph$$

the pullback to Σ .

The isometric embedding $\aleph$ also induces an isometry between the (intrinsic) spinor bundles of Σ_{i_0} and $\check{\Sigma}_{i_0}$ together with their Dirac operators which are isomorphic to $e_r \cdot D$ and $\check{e}_r \cdot \check{D}$ respectively. This isometry can be extended to an isometry over the complex 2-dimensional sub-bundles of their hypersurface spinor bundles. Denote by $\check{\mathbb{S}}^{\check{\Sigma}_{i_0}}$ this sub-bundle of $\check{\mathbb{S}}|_{\check{\Sigma}_{i_0}}$. Let $\check{\phi}$ be a constant section of $\check{\mathbb{S}}^{\check{\Sigma}_{i_0}}$ and denote

$$\phi_0 = \check{\phi} \circ \aleph.$$

Denote by $\check{\Xi}$ the set of all these constant spinors $\check{\phi}$ with the unit norm. This set is isometric to S^3 .

Suppose that one of the following conditions holds on M

(i) $tr_g(p) \geq 0$, $H|_{\Sigma_i} + tr(p|_{\Sigma_i}) \geq 0$ for all i ;

(ii) $tr_g(p) \leq 0$, $H|_{\Sigma_i} - tr(p|_{\Sigma_i}) \geq 0$ for all i .

Let ϕ be the unique solution of (2.10) or (2.11) for some $\check{\phi} \in \check{\Xi}$. Denote

$$\begin{aligned} m(\Sigma_{i_0}, \check{\phi}) &= \frac{1}{8\pi} \Re \int_{\Sigma_{i_0}} \left[(H_0 - H)|\phi|^2 \right. \\ &\quad \left. + tr(p|_{\Sigma_{i_0}}) \langle \phi, e_0 \cdot e_r \cdot \phi \rangle \right. \\ &\quad \left. - p_{ar} \langle \phi, e_0 \cdot e_a \cdot \phi \rangle \right]. \end{aligned} \quad (3.1)$$

The *quasi local mass* of Σ_{i_0} is defined as

$$m(\Sigma_{i_0}) = \min_{\check{\Xi}} m(\Sigma_{i_0}, \check{\phi}). \quad (3.2)$$

If all Σ_i can be smoothly isometrically embedded into $\mathbb{R}^3$ in the Minkowski spacetime $\mathbb{R}^{3,1}$, we define the *quasi local mass* of Σ as

$$m(\Sigma) = \sum_i m(\Sigma_i). \quad (3.3)$$

It is obvious that $m(\Sigma_{i_0})$ vanishes if Σ_{i_0} is an embedding 2-sphere in $\mathbb{R}^3$. Let E, P_1, P_2, P_3 be the ADM energy-momentum of an asymptotically flat spacetime [19, 21, 16, 23]. Take Σ_{i_0} to be a sphere in an end. Then

$$\begin{aligned} \lim_{r \rightarrow \infty} m(\Sigma_{i_0}) &= \min_{\check{\Xi}} \langle \check{\phi}, E\check{\phi} + P_k dx_0 \cdot dx_k \cdot \check{\phi} \rangle \\ &= E - |P|. \end{aligned}$$

Theorem 1. *Let $(N, \tilde{g})$ be a spacetime which satisfies the dominant energy condition. Let (M, g, p) be a smooth initial data set with the boundary Σ which has finitely many multi-components Σ_i , each of which is topological 2-sphere. Suppose that some Σ_{i_0} can be smoothly isometrically embedded into $\mathbb{R}^3$ whose image in $\mathbb{R}^3$ has nonnegative mean curvature. If either condition (i) or condition (ii) holds, then*

- (1) $m(\Sigma_{i_0}) \geq 0$;
- (2) that $m(\Sigma_{i_0}) = 0$ implies the energy-momentum of spacetime satisfies

$$T_{00} = |f||\phi|^2, \quad T_{0i} = f \langle \phi, e_0 \cdot e_i \cdot \phi \rangle$$

along M , where f is a real function, ϕ is the unique solution of (2.10) or (2.11) for some $\check{\phi} \in \check{\Xi}$.

- (3) Furthermore, if $p_{ij} = 0$, then $m(\Sigma_{i_0}) = 0$ implies that M is flat with connected boundary.

Proof : Note that (2.8) implies

$$\check{\nabla}_a \check{\phi} + \frac{1}{2} \check{h}_{ab} \check{e}_r \cdot \check{e}_b \cdot \check{\phi} = 0$$

on $\check{\Sigma}$. Then

$$e_r \cdot D\phi_0 = \frac{H_0}{2} \phi_0$$

on Σ . Denote $\phi_0^\pm = P_\pm \phi_0$. Since $e_r \cdot D \circ P_\pm = P_\mp \circ e_r \cdot D$, we have

$$e_r \cdot D\phi_0^\pm = \frac{H_0}{2} \phi_0^\mp.$$

Therefore, using the self-adjointness of $e_r \cdot D$, we obtain

$$\int_\Sigma H_0 |\phi_0^+|^2 = \int_\Sigma H_0 |\phi_0^-|^2.$$

Denote $\phi^\pm = P_\pm \phi$. Then $\int_\Sigma \langle \phi^+, \phi^- \rangle = 0$. In case (i), we have $\phi^+ = \phi_0^+$. Thus

$$\begin{aligned} \int_\Sigma \langle \phi, e_r \cdot D\phi \rangle &= 2\Re \int_\Sigma \langle \phi^-, e_r \cdot D\phi^+ \rangle \\ &= 2\Re \int_\Sigma \langle \phi^-, e_r \cdot D\phi_0^+ \rangle \\ &= \Re \int_\Sigma \langle \phi^-, H_0 \phi_0^- \rangle \\ &\leq \frac{1}{2} \int_\Sigma H_0 (|\phi^-|^2 + |\phi_0^-|^2) \\ &= \frac{1}{2} \int_\Sigma H_0 (|\phi^-|^2 + |\phi_0^+|^2) \\ &= \frac{1}{2} \int_\Sigma H_0 |\phi|^2. \end{aligned}$$

Therefore $m(\Sigma_{i_0}, \check{\phi}) \geq 0$ which implies the positivity. Same argument applies to the case (ii). Thus the proof of the first part is complete.

For the second part, if $m(\Sigma_{i_0}) = 0$, then there exists $\check{\phi}$ such that $m(\Sigma_{i_0}, \check{\phi}) = 0$. Let ϕ be the unique solution of (2.10) or (2.11) corresponding to the boundary value $\check{\phi}$. Then (2.9) implies $\check{\nabla}_i \phi = 0$ on M . This gives that $|\phi|$ is nonzero and

$$\langle \phi, (T_{00} + T_{0i} e_0 \cdot e_i) \phi \rangle = 0$$

along M . Therefore, the dominant energy condition and Lemma 1 [11] solve T_{00} and T_{0i} . Furthermore, if $p_{ij} = 0$ in this case, then there exists a parallel spinor on M . Thus M is Ricci flat, hence flat since it is

3-dimensional. As each component of its boundary has nonnegative mean curvature, the boundary must be connected. Q.E.D.

Remark 1. *The vanishing quasi local mass in Theorem 1 gives rise to one spinor which is parallel along M and it does not imply the Ricci flatness of the spacetime along M . This is similar to Dougan-Mason's quasi local mass [7]. There is a standard model for 4-dimensional spacetimes, i.e., pp -manifolds, admitting one parallel spinor,*

$$\tilde{g}_f = -2dx_1dx_2 + f(x_2, x_3, x_4)dx_2^2 + dx_3^2 + dx_4^2,$$

where f is a smooth function (e.g., [1]). $\tilde{g}_f$ is Ricci flat if and only if f satisfies

$$\frac{\partial^2 f}{\partial x_3^2} + \frac{\partial^2 f}{\partial x_4^2} = 0.$$

Remark 2. *The second fundamental form of the time slice in the Kerr spacetime is nontrivial and falls off as $\frac{1}{r^3}$ (e.g. [24]). Thus, for the regular initial data set which is Kerr at infinity constructed in [5], the “quasi-local momentum” of a bounded domain with sufficiently large radius may be nontrivial although the total ADM linear momentum is zero.*

We shall use null normals to express $m(\Sigma_{i_0}, \check{\phi})$. At each point of Σ_{i_0} , we choose two null normal vectors n_+ and n_- such that $\tilde{g}(n_+, n_-) = -1$. We also choose the local frame e_2, e_3 of Σ . Denote by $\mathbf{H} \equiv -\tilde{\nabla}_{e_2}e_2 - \tilde{\nabla}_{e_3}e_3$ the mean curvature vector of Σ . Let

$$2\rho = \tilde{g}(\mathbf{H}, n_+), \quad 2\mu = \tilde{g}(\mathbf{H}, n_-), \quad \varpi_a = -\tilde{g}(\tilde{\nabla}_a n_+, n_-)$$

where $a = 2, 3$. A straightforward computation yields

$$\text{tr}(p|_{\Sigma_{i_0}}) = \sqrt{2}(\rho - \mu), \quad p_{ar} = \varpi_a.$$

Thus

$$\begin{aligned} m(\Sigma_{i_0}, \check{\phi}) &= \frac{1}{8\pi} \Re \int_{\Sigma_{i_0}} \left[(H_0 - H)|\phi|^2 \right. \\ &\quad - \frac{\rho - \mu}{\sqrt{2}} \langle \phi, (n_+ \cdot n_- \cdot -n_- \cdot n_+ \cdot) \phi \rangle \\ &\quad \left. - \frac{\varpi_a}{\sqrt{2}} \langle \phi, (n_+ + n_-) \cdot e_a \cdot \phi \rangle \right]. \end{aligned}$$

This formula suggests a way to define the quasi-local mass for space-like 2-spheres Σ using geometric data of their embedding into a 4-dimensional spacetime. However, it is still unclear what the correct choice of spinor ϕ is. In the cases that Σ is a past apparent horizon and encloses a spacelike hypersurface M^+ with nonnegative mean

curvature, or Σ is a future apparent horizon and encloses a space-like hypersurface M^- with nonpositive mean curvature, then we can choose spinor ϕ as the solution (2.10) or (2.11). Denote $m(\Sigma_{i_0}, \check{\phi})$ as $m(\Sigma, \check{\phi}, M^\pm)$ for a chosen spacelike hypersurface $M^\pm$. Then we can define the quasi-local mass of Σ as

$$m(\Sigma) = \inf_{\{M^\pm\}} \min_{\check{\phi}} m(\Sigma, \check{\phi}, M^\pm)$$

where $\{M^\pm\}$ is the set of all required spacelike hypersurfaces enclosed by Σ . If Σ satisfies all conditions mentioned above, we can still prove $m(\Sigma)$ is nonnegative.

4. SCHWARZSCHILD SPACETIME

We conjecture that $m(\Sigma)$ has increasing monotonicity. We verify that it is indeed the case for the round spheres in time-slices M of the exterior Schwarzschild spacetime. For this M , the metric is

$$g_{Sch} = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\psi^2)$$

($r \geq 2m$) and the second fundamental form vanish, i.e., $p_{ij} = 0$. Using the isotropic coordinate $r = \rho\left(1 + \frac{m}{2\rho}\right)^2$ ($\rho \geq \frac{m}{2}$), g_{Sch} can be written as

$$g_{Sch} = \left(1 + \frac{m}{2\rho}\right)^4 [d\rho^2 + \rho^2(d\theta^2 + \sin^2\theta d\psi^2)] \equiv u^4 g_{flat}$$

where $u = 1 + \frac{m}{2\rho}$. g_{Sch} has an apparent horizon $\Sigma = \{\rho = \frac{m}{2}\}$ which is a minimal surface.

We briefly discuss the Dirac operator for conformal metrics. We refer to [10] and references therein for detail. Denote by G_u the isometry between $SO_{g_{flat}}$ and $SO_{g_{Sch}}$ given by the above conformal change of metric. The isometry G_u induces an isometry between the $Spin_3$ principal bundles as well as an isometry between their spinor bundles S and $\tilde{S}$ ($\equiv G_u S$). For any sections ϕ, ψ of the spinor bundle S , we denote $\tilde{\phi} = G_u \phi$, and $\tilde{\psi} = G_u \psi$ the corresponding sections of $\tilde{S}$. If $(\cdot, \cdot)_{g_{flat}}$ and $(\cdot, \cdot)_{g_{Sch}}$ denote respectively the natural Hermitian metrics on S and $\tilde{S}$, then

$$(\phi, \psi)_{g_{flat}} = (\tilde{\phi}, \tilde{\psi})_{g_{Sch}}.$$

The Clifford multiplication on $\tilde{S}$ is given by

$$\tilde{e}_i \cdot \tilde{\phi} = \widetilde{e_i \cdot \phi}.$$

Dirac operators with respect to the two metric have the relation

$$\overline{D}_{Sch}(u^{-1}\tilde{\phi}) = u^{-3}\widetilde{\overline{D}_{flat}\phi}.$$

For any covariant constant spinor $\check{\phi}$ with unit norm, $u^{-1}\tilde{\phi}$ is a solution of $\overline{D}_{Sch}$. Moreover, $|\tilde{\phi}|_{Sch}^2 = |\check{\phi}|_{flat}^2 = 1$.

For any round sphere B_ρ in M with radius ρ , the Brown-York quasi-local mass is (e.g. [14])

$$\begin{aligned} m_{BY}(B_\rho) &= \frac{1}{8\pi} \int_{S_\rho} (H_0 - H) \\ &= r \left(1 - \sqrt{1 - \frac{2m}{r}} \right) \\ &= m \left(1 + \frac{m}{2\rho} \right). \end{aligned}$$

Since $H_0 > 0$, the new quasi-local mass is

$$\begin{aligned} m(B_\rho) &= \min_{\tilde{\phi}} \frac{1}{8\pi} \int_{S_\rho} (H_0 - H) |u^{-1}\tilde{\phi}|_{Sch}^2 \\ &= r \left(1 - \sqrt{1 - \frac{2m}{r}} \right) u^{-2} \\ &= \frac{m}{1 + \frac{m}{2\rho}}. \end{aligned}$$

Therefore $m(B_\rho)$ is increasing and

$$\lim_{\rho \rightarrow \infty} m(B_\rho) = m = E_{ADM}.$$

Acknowledgements. The author is indebted to M.T.Wang and S.T.Yau for sharing their idea and some valuable conversations.

REFERENCES

- [1] H. Baum, *Twistor and Killing spinors in Lorentzian geometry*, Global analysis and harmonic analysis (Marseille-Luminy, 1999), 35-52, Semin. Congr. 4, Soc. Math. France, Paris, 2004.
- [2] J.D. Brown and J.W. York, *Quasilocal energy in general relativity*, Mathematical aspects of classical field theory (Seattle, WA, 1991), 129-142, Contemp. Math., 132, Amer. Math. Soc., Providence, RI, 1992.
- [3] J.D. Brown and J.W. York, *Quasilocal energy and conserved charges derived from the gravitational action*, Phys. Rev. D(3) 47, 1407-1419 (1993).
- [4] J.D. Brown, S.R. Lau and J.W. York, *Energy of isolated systems at retarded times as the null limit of quasilocal energy*, Phys. Rev. D 55, 1977-1984 (1997).

- [5] J. Corvino and R. Schoen, *On the asymptotics for the vacuum Einstein constraint equations*, J. Diff. Geom. 73, 185-217 (2006).
- [6] D. Christodoulou and S.T. Yau, *Some remarks on the quasi-local mass*, Mathematics and general relativity, 9-14, Contemp. Math., 71, Amer. Math. Soc., Providence, RI, 1988.
- [7] A.J. Dougan and L.J. Mason, *Quasilocal mass constructions with positive energy*, Phys. Rev. Lett. 67, 2119-2122 (1991).
- [8] G.W. Gibbons, S.W. Hawking, G.T. Horowitz and M.J. Perry, *Positive mass theorems for black holes*, Commun. Math. Phys. 88, 295-308 (1983).
- [9] S.W. Hawking and G.T. Horowitz, *The gravitational Hamiltonian, action, entropy and surface terms*, Classical Quantum Gravity 13, 1487-1498 (1996).
- [10] O. Hijazi and X. Zhang, *Lower bounds for eigenvalues of the Dirac operator. Part I. The hypersurface Dirac operator*, Ann. Global Anal. Geom. 19, 355-376 (2001).
- [11] O. Hijazi and X. Zhang, *The Dirac-Witten operator on spacelike hypersurfaces*, Comm. Anal. Geom. 11, 737-750 (2003).
- [12] J. Kijowski, *A simple derivation of canonical structure and quasi-local Hamiltonians in general relativity*, Gen. Relat. Grav. 29, 307-343 (1997).
- [13] C-C.M. Liu and S.T. Yau, *Positivity of quasilocal mass*, Phys. Rev. Lett. 90, 231102 (2003).
- [14] C-C.M. Liu and S.T. Yau, *Positivity of quasilocal mass II*, J. Amer. Math. Soc. 19, 181 (2006).
- [15] N.Ó Murchadha, L.B. Szabados and K.P. Tod, *Comment on "Positivity of quasilocal mass"*, Phys. Rev. Lett. 92, 259001 (2004).
- [16] T. Parker and C.H. Taubes, *On Witten's proof of the positive energy theorem*, Commun. Math. Phys. 84, 223-238 (1982).
- [17] Y. Shi and L-F. Tam, *Positive mass theorem and the boundary behavior of compact manifolds with nonnegative scalar curvature*, J. Diff. Geom. 62, 79 (2002).
- [18] R. Schoen, S.T. Yau, *On the proof of the positive mass conjecture in general relativity*, Commun. Math. Phys. 65, 45-76 (1979).
- [19] R. Schoen, S.T. Yau, *Proof of the positive mass theorem. II*, Commun. Math. Phys. 79, 231-260 (1981).
- [20] M.T. Wang and S.T. Yau, *A generalization of Liu-Yau's quasi-local mass*, math.DG/0602321.
- [21] E. Witten, *A new proof of the positive energy theorem*, Commun. Math. Phys. 80, 381-402 (1981).
- [22] S.T. Yau *Geometry of three manifolds and existence of black hole due to boundary effect*, Adv. Theor. Math. Phys. 5, 755-767 (2001).
- [23] X. Zhang, *Positive mass conjecture for 5-dimensional Lorentzian manifolds*, J. Math. Phys. 40, 3540-3552 (1999).
- [24] X. Zhang, *Remarks on the total angular momentum in general relativity*, Commun. Theor. Phys. 39, 521-524 (2003).

INSTITUTE OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS
SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100080, CHINA

E-mail address: xzhang@amss.ac.cn